



After Epilation Fluid, Type O/W with AGENAFLO 9050

GUIDE FORMULATION

| Ingredients | INCI-Name | %w/w | Supplier |
|----------------------|---|------------|--------------------|
| Phase A | | | |
| Water | Aqua | Ad 100 | |
| Keltrol CG-SFT | Xanthan Gum | 0,2 | CP Kelco |
| Polyglycol 400 | Peg-8 | 3,0 | Clariant |
| Phase B | | | |
| AGENAFLO 9050 | Corn Starch Modified | 1,0 | AGRANA |
| Capigel 98 | Acrylates Copolymer | 2,0 | Seppic |
| Phase C | | | |
| Dow Corning 344 EU | Cyclomethicone | 3,0 | Dow Corning |
| Lactokine™ Fluid | Milk Protein | 4,0 | CLR |
| Hygroplex WHN | Hexylene Glycol, Fructose, Glucose, Sucrose, Urea, Dextrin, Alanine, Glutamic Acid, Aspartic Acid | 2,5 | CLR |
| Phase D | | | |
| Euxyl PE 9010 | Phenoxyethanol, Ethylhexylglycerin | 0,6 | Schuelke & Mayr |
| Phase E | | | |
| Aristoflex AVC | Ammonium Acryloyldimethyltaurate/VP Copolymer | 0,75 | Clariant |
| Phase F | | | |
| Alcohol Denat. | Alcohol Denat. | 5,0 | |

PROCEDURE

1. Mix A until completely soaked.
2. Add the other ingredients in the given order.
3. Stir until uniform.

This formulation is indicative only and no responsibility whatsoever will be assumed. This formulation is offered solely for your consideration, investigation and verification. It is your responsibility to thoroughly test any formulations before use. All those who use our formulations as well as those who process AGRANA Starches are themselves entirely responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies. AGRANA does not assure any liability with respect to the composition of the formulation or the use in products.